arXiv:1405.1017v3 [math.AP] 16 Mar 2016

A FOUNDATIONAL APPROACH TO THE LIE THEORY FOR FRACTIONAL ORDER PARTIAL DIFFERENTIAL EQUATIONS

ROSARIO ANTONIO LEO, GABRIELE SICURO, AND PIERGIULIO TEMPESTA

ABSTRACT. We provide a general theoretical framework allowing to extend the classical Lie theory for partial differential equations to the case of equations of fractional order. We propose a general prolongation formula allowing the study of Lie symmetries for case of an arbitrary finite number of independent variables. We also prove the Lie theorem in the case of fractional differential equations, showing that the proper space for the analysis of these symmetries is the infinite dimensional jet space.

CONTENTS

1. Introduction 1
2. Fundamental definitions 2
3. Lie Theory for Fractional PDEs 3
3.1. Local Lie groups 4
3.2. Jet space 4
3.3. General prolongation formula 6
3.4. The Lie theorem and its application 8
3.5. Reduction by symmetry 9
4. A nonlinear equation describing a process of anomalous diffusion 10
Acknowledgments 12
References 12

1. INTRODUCTION

The purpose of this paper is to provide a general analytic framework for the construction of exact solutions of fractional partial differential equations (FPDEs). The study of such equations has an intriguing history, dating back to XIX century [12, 6], with the works of Abel in classical mechanics, Riemann, Liouville, etc. In the last decade, there was a resurgence of interest in FDEs, since these equations have been recognized as fundamental not only in pure mathematics, but also in statistical mechanics, economics, social sciences, for their manifold applications especially in the description of nonlinear phenomena like anomalous diffusion. Consequently, the search for exact solutions of both stationary and evolution FPDEs is of great relevance. Although many analytic approaches have been largely (but not exhaustively) investigated, the determination of algebraic and geometric methods for the study of FPDEs is essentially an open problem.

Concerning group-theoretical methods, in [2] the invariance of a fractional diffusion equation under scaling transformations was first noticed. Subsequently, in [3, 4, 8] an extension of the Lie approach for standard differential equations to the case of FPDEs was proposed. However, in these works, only the case of equations

Date: March 17, 2016.

involving fractional derivatives with respect to a single independent variable has been analyzed, keeping the other ones as integer order derivatives.

Nowadays, several purely computational studies about symmetries of fractional order differential equations can be found in the literature. Nevertheless, to our knowledge a complete and rigorous mathematical theory is essentially lacking.

The present paper contributes to fill this gap. Our main result is a coherent approach to the notion of symmetries for FPDEs, which significantly generalize the known theoretical results (see in particular [3, 4]). We will show the existence of a natural connection between the theory of Lie invariance of FPDEs and the classical one for PDEs, once extended to the case of equations involving an infinite number of derivatives [14]. Our motivation is the fact that, in the case of standard PDEs, the theory of Lie symmetries proved to be one of the most effective tools for the determination of classes of analytic solutions [10, 1, 5, 15]. Since the crucial observation, due to G. Birchoff, of the potential relevance of Lie theory in the study of the equations of fluid dynamics, in the middle of the XX century, an impressive body of literature has been produced on the applications of symmetries to both linear and nonlinear problems, which is still a flourishing research area.

Our extension of group–theoretical methods to the case of FPDEs comes from the geometrical notions of infinite prolongation and infinite dimensional jet space $\mathcal{J}^\infty$ [7, 13, 14]. Indeed, a FPDE can also be interpreted as a system of PDEs involving an infinite number of (classical) derivatives. Our main results are the following. First, we prove a generalized prolongation formula for the action of a vector field on fractional derivative operators involving an arbitrary number of independent variables. In the process, we recover as special cases the results already known in the literature. Second, by means of this prolongation formula, we derive the Lie condition for a vector field to be an infinitesimal generator of a symmetry group.

As in the classical approach, the vector fields defined by the previous condition possess a twofold role. First, they generate diffeomorphisms that foliate the manifold of the independent and dependent variables into group orbits, transforming solutions into solutions. Second, and more importantly, they allow to reduce a given FPDE into another one, possessing fewer independent variables, or a fractional ordinary differential equation (FODE) into an equation of reduced order. Solving the reduced equations enables us to construct exact solutions of the original problem.

The results obtained in this paper pave the way to the creation of a complete theory of Lie symmetries for FPDEs, that would include generalizations to higher order symmetries, nonclassical symmetries, master symmetries, etc.

The paper is organized as follows. In Section 2, some basic definitions concerning the fractional calculus are reviewed. Section 3 is devoted to the notion of symmetry for FPDEs: we show the correspondence between the fractional theory and the classical one and we derive explicitly a generalized prolongation formula expressing the action of the infinitely prolonged symmetry field on fractional operators. In Section 4, the general prolongation formula of a vector field in a suitable space is obtained. Finally, an application of our approach to a $(N+1)$-dimensional nonlinear diffusive FPDE is presented.

2. Fundamental definitions

In this section, we will review briefly some basic definitions of fractional calculus. For an extensive and detailed treatment of the subject and the proofs of the results briefly presented here, we refer to [12, 6]. Given $p \in \mathbb{R}$, we introduce the symbol

$$\langle p \rangle := \begin{cases} [p] & p \geq 0, \\ -1 & p < 0 \end{cases} \tag{2.1}$$

In the previous expression $[p] \in \mathbb{Z}$ is the floor function of p, $[p] \leq p < [p] + 1$. We can now state the following

Definition 2.1 (Riemann–Liouville fractional operator)**.** Let $p \in \mathbb{R}$ and $f\colon [a,b] \subset \mathbb{R} \to \mathbb{R}$ be a real analytic function in (a,b). We shall call

$$ {}_a\mathcal{D}_t^p f(t) \coloneqq \frac{1}{\Gamma(1+\langle p\rangle - p)} \frac{\mathrm{d}^{\langle p\rangle+1}}{\mathrm{d}\, t^{\langle p\rangle+1}} \int_a^t (t-\tau)^{\langle p\rangle - p} f(\tau)\, \mathrm{d}\,\tau, \qquad t \in (a,b), \tag{2.2} $$

the Riemann–Liouville (RL) fractional operator of order p and terminals (a,t). Here $\Gamma(x) \coloneqq \int_0^\infty t^{x-1}\, \mathrm{e}^{-t}\, \mathrm{d}\, t$ is the Gamma function.

The operator defined above is usually called the *RL fractional derivative* when $p > 0$ and the *RL fractional integral* for $p < 0$. Observe that the RL operator is linear and for $k \in \mathbb{Z}$,

$$ \lim_{\substack{p\to k^{\pm}\\ k\in\mathbb{Z}}} {}_a\mathcal{D}_t^p f(t) = \begin{cases} \frac{\mathrm{d}^k f(t)}{\mathrm{d}\, t^k} & \text{if } k > 0,\\ f(t) & \text{if } k = 0,\\ \frac{1}{(|k|-1)!} \int_a^t (t-\tau)^{|k|-1} f(\tau)\, \mathrm{d}\,\tau & \text{if } k < 0. \end{cases} \tag{2.3} $$

Similarly, we can introduce the partial RL operator as follows.

Definition 2.2. Let $p_i \in \mathbb{R}$, $i = 1, \dots, N$, and

$$ f(x_1, \dots, x_N)\colon [a_1, b_1] \times \cdots \times [a_N, b_N] \subset \mathbb{R}^N \to \mathbb{R}, $$

be a real analytic function in $(a_1, b_1) \times \cdots \times (a_N, b_N)$. We define the mixed RL operator as [6]

$$ \begin{aligned} {}_{a_1,\dots,a_N}\partial^{p_1,\dots,p_N}_{x_1,\dots,x_N} f(x_1,\dots,x_N) \coloneqq \\ = \left(\prod_{i=1}^N \frac{\partial^{\langle p_i\rangle+1}}{\partial x_i^{\langle p_i\rangle+1}} \int_{a_i}^{x_i} \mathrm{d}\,\xi_i\, \frac{(x_i-\xi_i)^{\langle p_i\rangle - p_i}}{\Gamma(1+\langle p_i\rangle - p_i)} \right) f(\xi_1,\dots,\xi_N). \end{aligned} \tag{2.4} $$

Remarkably, a mixed RL operator acting on an analytic function can be formally written as

$$ {}_{a_1,\dots,a_N}\partial^{p_1,\dots,p_N}_{x_1,\dots,x_N} = \prod_{i=1}^N \left[\sum_{k_i=0}^\infty \binom{p_i}{k_i} \frac{(x_i-a_i)^{k_i-p_i}}{\Gamma(k_i-p_i+1)} \frac{\partial^{k_i}}{\partial x_i^{k_i}} \right]. \tag{2.5} $$

In the previous expression we have that

$$ \binom{p}{k} \coloneqq \frac{\prod_{i=1}^k (p-i+1)}{\Gamma(k+1)}. \tag{2.6} $$

Formula (2.5) will be fundamental to link the theory of symmetries of fractional differential equations with the classical theory of symmetries of PDEs. The relation has been proved in [9, Sec. 3.5] in the case of a fractional derivative operator with respect to one variable only acting on a space of analytic functions. The case of many variables can be treated similarly.

3. Lie Theory for Fractional Pdes

In this section we shall develop a theoretical framework allowing to generalize the classical Lie theory to the case of fractional partial differential equations.

3.1. **Local Lie groups.** We shall focus on *fractional partial differential equations*(FPDEs) with one dependent variable $u \in U \subseteq \mathbb{R}$ and N independent variables $(x_1, \dots, x_N) \in X := [a_1, +\infty) \times \cdots \times [a_N, +\infty) \subseteq \mathbb{R}^N$, i.e., on equations of the form

$$\mathcal{E}(x, u, {}_{\mathbf{a}}\partial^{\mathbf{p}}_x u, \dots) = 0, \tag{3.1}$$

where $\mathcal{E}$ is a function involving a finite number of RL operators

$${}_{\mathbf{a}}\partial^{\mathbf{p}}_x u := {}_{a_1,\dots,a_N}\partial^{p_1,\dots,p_N}_{x_1,\dots,x_N} u(x), \qquad p_i \in \mathbb{R}\, \forall i. \tag{3.2}$$

We will assume that all RL operators acting upon the same variable x_i have the same lower extreme a_i. By analogy with the case of PDEs we propose the following notion of symmetry group of a FPDE.

Definition 3.1 (Symmetries)**.** A *continuous group G of (point) symmetries* for the FPDE (3.1) is a local group of diffeomorphisms that map solutions $(x, u) \in M \subset X \times U =: \mathcal{M}$ (with M open subset of $\mathcal{M}$) into solutions $g \cdot (x, u) = (\tilde{x}, \tilde{u}) = (\Xi_g(x, u), \Phi_g(x, u)) \in X \times U$, $g \in G$, for some smooth functions $\Xi_g \colon M \to X$, $\Phi_g \colon M \to U$.

We denote by $\mathbf{v} \in \mathfrak{g}$ an element of the Lie algebra $\mathfrak{g}$ associated with the Lie group of transformations G. We shall consider vector fields of the form

$$\mathbf{v} = \sum_{i=1}^{N} \xi^i(x, u) \frac{\partial}{\partial x_i} + \phi(x, u) \frac{\partial}{\partial u}. \tag{3.3}$$

Here, assuming that the local action of the symmetry group G is given by

$$(x, u) \xrightarrow{g} (\tilde{x}, \tilde{u}), \quad g \equiv g(\varepsilon) = \mathrm{e}^{\varepsilon \mathbf{v}}, \quad \varepsilon \in \mathbb{R},$$

the following formulae hold:

$$\xi^i(x, u) = \left. \frac{\mathrm{d}\, \Xi^i_{g(\varepsilon)}(x, u)}{\mathrm{d}\, \varepsilon} \right|_{\varepsilon = 0}, \qquad \phi(x, u) = \left. \frac{\mathrm{d}\, \Phi_{g(\varepsilon)}(x, u)}{\mathrm{d}\, \varepsilon} \right|_{\varepsilon = 0}. \tag{3.4}$$

If a RL operator of u with respect to x_i appears in $\mathcal{E}$, we impose the additional constraint $a_i \xrightarrow{g} a_i$, i.e., $\xi^i(x_1, \dots, x_{i-1}, a_i, x_{i+1}, \dots x_N) \equiv 0$, i.e., ${}_{\mathbf{a}}\partial^{\mathbf{p}}_x u(x) \xrightarrow{g} {}_{\mathbf{a}}\partial^{\mathbf{p}}_{\tilde{x}} \tilde{u}(\tilde{x})$. To evaluate the functions (3.4), we need to *prolong* our function u in a suitable extended space, as we will show below.

3.2. **Jet space.** To treat properly the problem of symmetries of FPDEs, we need to implement an "infinite" prolongation procedure of vector fields over a suitable infinite-dimensional jet space $\mathcal{J}^\infty$. The theory of infinite dimensional jet bundles [13, 14, 7] is a generalization of the classical finite dimensional theory. In the following, we will use the standard multi-index notation for derivatives of a function $f(x) \equiv f(x_1, \dots, x_N)$, i.e., for a set of indices $\sigma = (i_1, \dots, i_K)$, $1 \leq i_k \leq N$, $k = 1, \dots, K$, we shall denote by

$$\frac{\partial^{|\sigma|} f(x)}{\partial x^\sigma} = \left(\prod_{k=1}^{K} \frac{\partial}{\partial x_{i_k}} \right) f(x), \quad \text{with } |\sigma| := K.$$

Definition 3.2 (Infinite jet manifold)**.** Let $(\mathcal{M}, \pi, \mathcal{X})$ a fiber bundle with projection $\pi \colon \mathcal{M} \to \mathcal{X}$ from the total space $\mathcal{M}$ to the base space $\mathcal{X}$. We assume $\dim \mathcal{M} = \dim \mathcal{X} + 1 = N + 1$. Then, denoting by $\Gamma_\eta(\pi)$ the set of all local sections of π containing $\eta \in \mathcal{X}$, we say that two sections $f, \tilde{f} \in \Gamma_\eta(\pi)$ are ∞-*equivalent* in x if

$$\left. \frac{\partial^{|\sigma|} f(x)}{\partial x^\sigma} \right|_{x=\eta} = \left. \frac{\partial^{|\sigma|} \tilde{f}(x)}{\partial x^\sigma} \right|_{x=\eta} \tag{3.5}$$

for $0 \leq |\sigma| < \infty$. We call the equivalence class of f the ∞-*jet* of f in η, and we denote it by $j^\infty_\eta(f)$. The infinite jet manifold $\mathcal{J}^\infty(\pi)$ is the set of these equivalence classes

$$\mathcal{J}^\infty(\pi) \coloneqq \{j^\infty_\eta(f) \colon \eta \in \mathcal{X},\ f \in \Gamma_\eta(\pi)\}. \tag{3.6}$$

Given a fractional differential equation in the form of Eq. (3.1), the space introduced above represents the suitable space for the prolongation procedure we require, with the identifications $\mathcal{M} \equiv X \times U$ and $\mathcal{X} \equiv X$. Indeed, using Eq. (2.5) we can rewrite Eq. (3.1) as a differential equation involving an *infinite* number of ordinary partial derivatives. Consequently, Eq. (3.1) defines an infinite dimensional submanifold

$$\mathcal{L}^\infty \coloneqq \left\{j^\infty_\eta(f) \in \mathcal{J}^\infty(\pi) \colon \mathcal{E}(\eta, f, {}_{\mathbf{a}}\partial^{\mathbf{P}}_\eta f, \dots) = 0\right\} \subset \mathcal{J}^\infty(\pi). \tag{3.7}$$

An infinitesimal symmetry of the equation $\mathcal{E}$ is therefore generated by a *vector field* tangent to $\mathcal{L}^\infty$ which preserves the contact structure of $\mathcal{L}^\infty$. Observe that in the infinite dimensional case we cannot associate, in general, a *flow* to $\mathbf{v}$ [14]: if this happens, we shall call $\mathbf{v}$ a *Lie field* (note that we deal with one dependent variable only). *We are interested in Lie fields whose structure in local coordinates is*

$$\mathbf{v} = \sum_{i=1}^N \xi^i(x,u)\frac{\partial}{\partial x_i} + \phi(x,u)\frac{\partial}{\partial u} + \sum_{|\sigma|\geq 1} \phi^\sigma(x, u, \dots, \partial^\kappa_x u, \dots)\frac{\partial^{|\sigma|}}{\partial u^\sigma}, \quad u^\sigma \coloneqq \frac{\partial^{|\sigma|} u}{\partial x^\sigma}, \tag{3.8}$$

where ξ^i, ϕ and ϕ^σ are analytic functions of their variables.

Introducing the *total derivative operator*

$$\mathrm{D}_i \coloneqq \frac{\partial}{\partial x_i} + \sum_{|\sigma|\geq 0} \frac{\partial u^\sigma}{\partial x_i}\frac{\partial}{\partial u^\sigma}, \tag{3.9}$$

it can be shown [14] that *every vector field associated to an infinitesimal symmetry transformation* can be written as

$$\mathrm{pr}^\infty \mathbf{v} = \sum_{i=1}^N \xi^i(x,u)\frac{\partial}{\partial x_i} + \phi(x,u)\frac{\partial}{\partial u} + \sum_{|\sigma|\geq 1} \phi^\sigma \frac{\partial}{\partial u^\sigma} \tag{3.10}$$

where, for $\sigma \equiv (i_1, \dots, i_K)$, $\mathrm{D}^\sigma \coloneqq \mathrm{D}_{i_1} \circ \cdots \circ \mathrm{D}_{i_K}$, and

$$\phi^\sigma \coloneqq \mathrm{D}^\sigma\left(\phi(x,u) - \sum_{i=1}^N \xi^i(x,u)\frac{\partial u}{\partial x_i}\right) + \sum_{i=1}^N \xi^i(x,u)\frac{\partial}{\partial x_i}\mathrm{D}^\sigma u. \tag{3.11}$$

The latter quantity depends on $N+1$ functions $\xi^1, \dots, \xi^N, \phi$ to be determined. We say that the vector field in Eq. (3.10) expresses the *prolonged action* of the vector field

$$\mathbf{v} = \sum_{i=1}^N \xi^i(x,u)\frac{\partial}{\partial x_i} + \phi(x,u)\frac{\partial}{\partial u} \in \mathsf{T}M. \tag{3.12}$$

The problem of local symmetries of a FPDE can be solved in the geometrical framework of the infinite dimensional jet space $\mathcal{J}^\infty$ of integer order derivatives introduced above; consequently, the theory of symmetry reduction is automatically borrowed from the classical theory. Indeed, from Eq. (2.5), we have that a fractional derivative can always be represented though a system of an infinite number of classical differential equations. For example, let us consider the fractional differential equation

$${}_a\mathcal{D}^p_x\, u(x) = f(x), \quad p \in \mathbb{R}. \tag{3.13}$$

Then, we can write

$$
{}_a\mathcal{D}_x^p\, u(x) = f(x) \Leftrightarrow \begin{cases} z_0 = f(x) \\ \omega_0 = u \\ \omega_k = \frac{\mathrm{d}\,\omega_{k-1}}{\mathrm{d}\,x}, & k \in \mathbb{N}, \\ z_k = z_{k-1} - \binom{p}{k-1} \frac{(x-a)^{k-p-1}}{\Gamma(k-p)} \omega_{k-1}, & k \in \mathbb{N}. \end{cases} \tag{3.14}
$$

A FPDE defines an infinite dimensional manifold in the space $\mathcal{J}^\infty$ endowed with an infinite order contact structure. The problem of finding a symmetry for a FPDE (i.e., the corresponding functions $\xi^1, \dots, \xi^N, \phi$) therefore can be formulated on the same geometric structures introduced in the classical theory of higher symmetries [14]. It will be solved in the space $\mathcal{J}^\infty$ by imposing the contact condition provided by a suitable version of the standard Lie theorem, that will be discussed below.

3.3. **General prolongation formula.** We look now for an explicit expression for the quantity

$$
\phi^{\mathbf{p}} \coloneqq \mathrm{pr}^\infty \mathbf{v}\, [\, {}_{\mathbf{a}}\partial_x^{\mathbf{p}}\, u]\,, \tag{3.15}
$$

as a function of the coefficients of $\mathbf{v}$. To stress the formal analogy with Eq. (3.11), we call the expression for $\phi^{\mathbf{p}}$ the *general prolongation formula*.

Theorem 3.1 (General prolongation formula)**.** *Let $\mathbf{v} = \sum_{i=1}^N \xi^i(x,u)\partial_{x_i} + \phi(x,u)\partial_u$ be a vector field defined on $\mathsf{T}M$, $M \subseteq \mathcal{M} = X \times U$, where X is an N–dimensional space, whose generic element has the form $x = (x_1, \dots, x_N) \in X$, and U is a one dimensional space. The general prolongation formula is given by*

$$
\phi^{\mathbf{p}} \coloneqq \mathrm{pr}^\infty \mathbf{v}\, [\, {}_{\mathbf{a}}\partial_x^{\mathbf{p}}\, u] = {}_{\mathbf{a}}\mathcal{D}_x^{\mathbf{p}} \left(\phi - \sum_{i=1}^N \xi^i \partial_{x_i} u \right) + \sum_{i=1}^N \xi^i \partial_{x_i}\, {}_{\mathbf{a}}\partial_x^{\mathbf{p}}\, u. \tag{3.16}
$$

In the previous equation we introduced the fractional total derivative

$$
{}_{\mathbf{a}}\mathcal{D}_x^{\mathbf{p}} \coloneqq \prod_{i=1}^N \left[\sum_{k_i=0}^\infty \binom{p_i}{k_i} \frac{(x_i - a_i)^{k_i - p_i}}{\Gamma(k_i - p_i + 1)}\, \mathrm{D}_i^{k_i} \right]. \tag{3.17}
$$

Note that the classical prolongation formula [1, 5, 10] is immediately recovered in the limit case of the integer order differentiation.

Proof. Using Eq. (2.5), we shall apply the vector field (3.10) to

$$
{}_{\mathbf{a}}\partial_x^{\mathbf{p}}\, u \coloneqq u \prod_{i=1}^N \frac{(x_i - a_i)^{-p_i}}{\Gamma(1-p_i)} + \sum_{|\sigma| \geq 1} C_\sigma(x) u^\sigma. \tag{3.18}
$$

In the latter expression, $\sigma = (\sigma_1, \dots, \sigma_K)$ is a multi-index and

$$
k_i(\sigma) \coloneqq \sum_{j=1}^K \delta_{\sigma_j, i}, \quad C_\sigma(x) \coloneqq \prod_{i=1}^N \frac{(x_i - a_i)^{k_i(\sigma) - p_i}}{\Gamma(k_i(\sigma) - p_i + 1)}. \tag{3.19}
$$

We obtain directly

$$
\begin{aligned}
(3.20)\quad \phi^{\mathbf{P}} &:= \mathrm{pr}^{\infty}\,\mathbf{v}\,[\,{}_{\mathbf{a}}\partial_x^{\mathbf{P}}\,u] = \\
&= \phi\prod_{i=1}^{N}\frac{(x_i-a_i)^{-p_i}}{\Gamma(1-p_i)} + \sum_{|\sigma|\geq 1} C_\sigma(x)\phi^\sigma + u\sum_{i=1}^{N}\xi^i\frac{(x_i-a_i)^{-p_i-1}}{\Gamma(-p_i)}\prod_{j\neq i}\frac{(x_j-a_j)^{-p_j}}{\Gamma(1-p_j)} \\
&\quad + \sum_{i=1}^{N}\xi^i\sum_{|\sigma|\geq 1}u^\sigma\frac{(x_i-a_i)^{k_i(\sigma)-p_i-1}}{\Gamma(k_j(\sigma)-p_j)}\prod_{j\neq i}^{N}\frac{(x_j-a_j)^{k_j(\sigma)-p_j}}{\Gamma(k_j(\sigma)-p_j+1)} \\
&\quad = {}_{\mathbf{a}}\mathcal{D}_x^{\mathbf{P}}\,\phi - \sum_{i=1}^{N}{}_{\mathbf{a}}\mathcal{D}_x^{\mathbf{P}}\left(\xi^i\partial_{x_i}u\right) + \sum_{i=1}^{N}\xi^i\partial_{x_i}\,{}_{\mathbf{a}}\partial_x^{\mathbf{P}}\,u,
\end{aligned}
$$

which coincides with the general prolongation formula (3.16). □

Corollary 3.2 (Prolongation formula for $N=1$)**.** *Under the hypotheses of Theorem 3.1, let us consider $N=1$ and $\mathbf{v}=\xi(x,u)\partial_x+\phi(x,u)\partial_u\in\mathsf{T}M$, $M\subset\mathcal{M}:=X\times U$. The general prolongation formula is given by*

$$
(3.21)\qquad \phi^p = {}_a\partial_x^p\,\phi - \sum_{n=0}^{\infty}\binom{p}{n+1}\frac{\mathrm{d}^{n+1}\xi}{\mathrm{d}\,x^{n+1}}\,{}_a\partial_x^{p-n}\,u.
$$

For $a=0$ and $p>0$, it is convenient to make explicit the dependence on u of the coefficient $\phi\equiv\phi(x,u(x))$ appearing in the vector field of eq. (3.21). To this purpose, we recall a formula due to Osler [11]

$$
\begin{aligned}
(3.22)\quad {}_0\mathcal{D}_t^p\,f(t,g(t)) &= \\
&= \sum_{n=0}^{\infty}\sum_{m=0}^{n}\sum_{k=0}^{m}\sum_{r=0}^{k}\binom{p}{n}\binom{n}{m}\binom{k}{r}\frac{t^{n-p}(-g)^r}{k!\Gamma(n+1-p)}\frac{\mathrm{d}^m\,g^{k-r}}{\mathrm{d}\,t^m}\frac{\partial^{n-m+k}f(t,g)}{\partial t^{n-m}\partial g^k}.
\end{aligned}
$$

Isolating the terms linear in u an its derivatives, we have

$$
(3.23)\qquad {}_0\mathcal{D}_x^p\,\phi(x,u(x)) = {}_0\partial_x^p\,\phi - u\,{}_0\partial_x^p\,(\partial_u\phi) + \partial_u\phi\,{}_0\partial_x^p\,u + \sum_{k=1}^{\infty}\binom{p}{k}\frac{\partial^{k+1}\phi}{\partial x^k\partial u}\,{}_0\partial_x^{p-k}\,u + \mu^p(x,u).
$$

Also, in eq. (3.23) we have denoted with $\mu^p(x,u)$ the expression

$$
(3.24)\qquad \mu^p := \sum_{n=2}^{\infty}\sum_{m=2}^{n}\sum_{k=2}^{m}\sum_{r=0}^{k-1}\binom{p}{n}\binom{n}{m}\binom{k}{r}\frac{x^{n-p}(-u)^r}{k!\Gamma(n+1-p)}\frac{\mathrm{d}^m}{\mathrm{d}\,x^m}\left(u^{k-r}\right)\frac{\partial^{n-m+k}\phi(x,u)}{\partial x^{n-m}\partial u^k}.
$$

We deduce that, for $p>0$

$$
\begin{aligned}
(3.25)\quad \phi^p &= {}_0\partial_x^p\,\phi + {}_0\partial_x^p\,u\left(\partial_u\phi - p\frac{\mathrm{d}\,\xi}{\mathrm{d}\,x}\right) - u\,{}_0\partial_x^p\,\partial_u\phi \\
&\quad + \mu^p(x,u) + \sum_{n=1}^{\infty}\left[\binom{p}{n}\partial_x^n\partial_u\phi - \binom{p}{n+1}\frac{\mathrm{d}^{n+1}\,\xi}{\mathrm{d}\,x^{n+1}}\right]{}_0\partial_x^{p-n}\,u.
\end{aligned}
$$

For further details we refer to [4].

Corollary 3.3. *Assume that the hypotheses of Theorem 3.1 hold with $N = 2$, $x_1 \equiv x$, $x_2 \equiv t$ and $a_2 = 0$. For $p > 0$, we have*

$$
\begin{aligned}
(3.26)\quad \phi_t^p \coloneqq \phi^{0,p} = {}_0\partial_t^p\,\phi + {}_0\partial_t^p\,u\left(\partial_u\phi - p\frac{\mathrm{d}\,\tau}{\mathrm{d}\,t}\right) - u\,{}_0\partial_t^p\,\partial_u\phi \\
+\sum_{n=1}^{\infty}\left\{\left[\binom{p}{n}\partial_t^n\partial_u\phi - \binom{p}{n+1}\frac{\mathrm{d}^{n+1}\tau}{\mathrm{d}\,t^{n+1}}\right]{}_0\partial_t^{p-n}\,u - \binom{p}{n}\frac{\mathrm{d}^n\,\xi}{\mathrm{d}\,t^n}\partial_x\,{}_0\partial_t^{p-n}\,u\right\} \\
+\sum_{n=2}^{\infty}\sum_{m=2}^{n}\sum_{k=2}^{m}\sum_{r=0}^{k-1}\binom{p}{n}\binom{n}{m}\binom{k}{r}\frac{t^{n-p}(-u)^r}{k!\Gamma(n+1-p)}\frac{\mathrm{d}^m}{\mathrm{d}\,t^m}\left(u^{k-r}\right)\frac{\partial^{n-m+k}\phi}{\partial t^{n-m}\partial u^k}.
\end{aligned}
$$

A similar expression is obtained for $\phi^{p,0} =: \phi_x^p$.

The prolongation formula (3.26) can be found in [8]; however, there a different definition for the mixed RL operator is adopted.

3.4. **The Lie theorem and its application.** We can now state the following fundamental result.

Theorem 3.4 (Lie theorem for FPDEs)**.** *Let $\mathcal{E}(x, u, {}_a\partial_x^p\,u, \dots) = 0$ be a FPDEs, defined in the open subset $M \subset \mathcal{M}$. If*

$$
(3.27)\qquad \mathrm{pr}^{\infty}\,\mathbf{v}\left[\mathcal{E}\left(x, u, {}_{\mathbf{a}}\partial_x^{\mathbf{p}}\,u, \dots\right)\right]\big|_{\mathcal{E}=0} = 0,
$$

then $\mathbf{v} = \sum_{i=1}^{N}\xi^i(x,u)\frac{\partial}{\partial x_i} + \phi(x,u)\frac{\partial}{\partial u}$ generates an infinitesimal symmetry of the equation $\mathcal{E} = 0$.

This theorem is a direct consequence of the Lie theorem stated in [14] for equations of infinite order, and applied to the case of a FPDE written as in Eq. (3.14) as a system of equations of infinite order. To apply the previous theorem to a FPDE, we must expand in principle every RL operator using Eq. (2.5) and then apply the Lie equation (3.27) to determine the coefficients ξ^i, ϕ of all infinitesimal symmetry vector fields: the Lie equation provides, in principle, an infinite number of differential equations for the $N + 1$ components of the field $\mathbf{v}$. *The symmetries of the general* FPDE *are therefore obtained by solving the classical Lie condition in the standard infinite dimensional jet space $\mathcal{J}^{\infty}$.*

The determining equation coming from the Lie theorem can be quite involved, due to the infinite number of independent components in $\mathcal{J}^{\infty}$. Moreover, according to the previous analysis, different fractional operators cannot be considered, strictly speaking, to be independent in the space $\mathcal{J}^{\infty}$ (this fact can be easily seen considering that the cardinality of the set of fractional operators is larger then the cardinality of the set of integer order derivatives). If we consider ${}_{\mathbf{a}}\partial_x^{\mathbf{p}}\,u$ as an "effective coordinate" in $\mathcal{J}^{\infty}$ we can still obtain symmetries for the initial differential problem; in particular, the expressions for the action of the symmetry field on RL operators can be formulated in a more compact way. This, in turn, allows us to write down in a simpler manner the Lie equation (3.27) and the general prolongation formula in Eq. (3.16). However, proceeding in this manner, we will find in general a *subgroup* of the complete group of Lie symmetries of the equation. The assumption of independence is strictly correct if the coefficients of a linear combination of fractional derivatives depend on x, u and a finite number of derivatives of u only. For example, let us suppose that we want to search for the solution of the following functional equation in $\mathcal{J}^{\infty}$:

$$
(3.28)\qquad \psi_1\left(x, u, \frac{\mathrm{d}\,u}{\mathrm{d}\,x}, \dots, \frac{\mathrm{d}^l\,u}{\mathrm{d}\,x^l}\right){}_a\mathcal{D}_x^p\,u(x) + \psi_2\left(x, u, \frac{\mathrm{d}\,u}{\mathrm{d}\,x}, \dots, \frac{\mathrm{d}^m\,u}{\mathrm{d}\,x^m}\right){}_a\mathcal{D}_x^q\,u(x) = 0.
$$

with $p, q \in \mathbb{R} \setminus \mathbb{N}_0$ and $l, m \in \mathbb{N}$. The equation above can be written as

$$\sum_{k=0}^{\infty} \left[\psi_1 \binom{p}{k} \frac{(x, u, \dots, \partial_x^l u)(x-a)^{k-p}}{\Gamma(k-p+1)} + \psi_2 \binom{q}{k} \frac{(x-a)^{k-q}}{\Gamma(k-q+1)} \right] \frac{\partial^k u}{\partial x^k} = 0. \tag{3.29}$$

This implies for $k \geq \max\{l, m\}$

$$\psi_1 \frac{\Gamma(p+1)(x-a)^{-p}}{\Gamma(p-k+1)\Gamma(k-p+1)} + \psi_2 \frac{\Gamma(q+1)(x-a)^{-q}}{\Gamma(q-k+1)\Gamma(k-q+1)} = 0. \tag{3.30}$$

The fact that the previous relation must hold for all value of $k \geq \max\{l, m\}$, implies that $\psi_1 = \psi_2 = 0$.

3.5. **Reduction by symmetry.** In the standard case of PDEs [10, 1, 14], given a local group G of transformations of a differential equation, the theory of symmetry reduction is formulated in terms of the quotient manifold $\mathcal{M}/G$ of the orbits of G on $\mathcal{M}$, and of the *projection map* $\pi\colon \mathcal{M} \to \mathcal{M}/G$. This map associates with every point $(x, u) \in \mathcal{M}$ its orbit under the action of G. In particular, the existence of the quotient manifold $\mathcal{M}/G$ and the projection map π is assured if G acts regularly with s dimensional orbits on the $(N+1)$ dimensional manifold $\mathcal{M}$. The map $\mathrm{d}\,\pi\colon\ T\mathcal{M}|_{(x,u)} \to T(\mathcal{M}/G)|_{\pi(x,u)}$ is onto and its kernel is $\mathfrak{g}|_{(x,u)} \coloneqq \{\mathbf{v}|_{(x,u)} : \mathbf{v} \in \mathfrak{g}\}$, where $\mathfrak{g}$ is the Lie algebra associated with G. In the context of FPDEs, the same philosophy applies. Therefore, following the standard theory, given a set of local coordinates, the algebra $\mathfrak{g}$ of G is generated by

$$\mathbf{v}_k \coloneqq \sum_{i=1}^{N} \xi_k^i(x, u)\partial_{x_i} + \phi_k(x, u)\partial_u, \quad k = 1, \dots, r. \tag{3.31}$$

If s is the dimension of the orbit of G through $\tilde{z} = (\tilde{x}, \tilde{u}) \in \mathcal{M}$, the transversality condition can be expressed locally as

$$\operatorname{rank}\left[\xi_k^i(\tilde{x}, \tilde{u})\right] = \operatorname{rank}\left[\xi_k^i(\tilde{x}, \tilde{u}), \phi^k(\tilde{x}, \tilde{u})\right] = s. \tag{3.32}$$

Assume now that G provides $N + 1 - s$ globally defined functionally independent invariants on $\mathcal{M}$. Let us call $v = \zeta(x, u)$ the new dependent variable and $y_i = \eta_i(x, u)$ the new $N - s$ independent variables. By construction,

$$J(x, u) \coloneqq \begin{pmatrix} \partial_j \eta^i & \partial_u \eta^i \\ \partial_j \zeta & \partial_u \zeta \end{pmatrix} \tag{3.33}$$

has rank $N + 1 - s$ everywhere. From eq. (3.32), it follows that in the case of a single fractional equation, the local transversality condition has the form

$$\operatorname{rank}\left[\begin{pmatrix} \partial_u \eta^i \\ \partial_u \zeta \end{pmatrix}\right] = 1. \tag{3.34}$$

If the previous condition is satisfied, the reduction can be locally performed by using Dini's theorem and writing u and $N - s$ independent variables $\tilde{x}$ as a function of the $N - s$ quantities y_i, of the s remaining independent variables $\hat{x}$ and v:

$$u = \alpha(\hat{x}, y, v), \qquad \tilde{x} = \beta(\hat{x}, y, v). \tag{3.35}$$

Finally, condition (3.34) guarantees that we can write u in the form (3.35); thus, the reduction can be performed.

4. A nonlinear equation describing a process of anomalous diffusion

To illustrate our approach, let us now consider the following equation:

$$\frac{\partial u}{\partial t} = \sum_{i=1}^{N} \alpha_i u^{\beta_i} \, {}_0\partial_{x_i}^{p_i} u, \qquad \alpha_i, \beta_i \in \mathbb{R} \setminus \{0\}, \quad p_i \in \mathbb{R}^+ \setminus \mathbb{N}. \tag{4.1}$$

It has the form of a fractional equation in which t has the role of a time coordinate, whilst x_i, $i = 1, \dots, N$, can be interpreted as space coordinates in an N–dimensional space. In this equation, there are no mixed operators and only one integro-differential operator appears for each variable. We will solve the problem of determining symmetries for this equation under the hypothesis that fractional derivatives of different order are independent. As specified above, this working ansatz is fullfilled when the coefficients of the fractional derivatives depend at most on x, u and a finite number of derivatives of u.

We search for a Lie field of the form

$$\mathbf{v} = \tau(t, x_1, \dots, x_N, u)\frac{\partial}{\partial t} + \sum_{i=1}^{N} \xi^i(t, x_1, \dots, x_N, u)\frac{\partial}{\partial x_i} + \phi(t, x_1, \dots, x_N, u)\frac{\partial}{\partial u}. \tag{4.2}$$

The determining equation reads

$$\left\{ \phi_t^1 - \sum_{i=1}^{N} \left[\alpha_i u^{\beta_i} \phi_{x_i}^{p_i} + \alpha_i \beta_i u^{\beta_i - 1} \phi \, {}_0\partial_{x_i}^{p_i} u \right] \right\} \Big|_{\partial_t u = \sum_{i=1}^{N} \alpha_i u^{\beta_i} \, {}_0\partial_{x_i}^{p_i} u} = 0, \tag{4.3}$$

where we have used Eq. (3.16); here

$$\begin{aligned}
\phi_{x_i}^p = {}& {}_0\partial_{x_i}^p \phi + {}_0\partial_{x_i}^p u \left(\frac{\partial \phi}{\partial u} - p \frac{\mathrm{d}\,\xi^i}{\mathrm{d}\,x_i} \right) - u \, {}_0\partial_{x_i}^p \frac{\partial \phi}{\partial u} - \sum_{n=1}^{\infty} \binom{p}{n} \frac{\mathrm{d}^n \tau}{\mathrm{d}\,x_i^n} \frac{\partial}{\partial t} \, {}_0\partial_{x_i}^{p-n} u \\
& + \sum_{n=1}^{\infty} \left\{ \left[\binom{p}{n} \frac{\partial^{n+1} \phi}{\partial x_i^n \partial u} - \binom{p}{n+1} \frac{\mathrm{d}^{n+1} \xi^i}{\mathrm{d}\,x_i^{n+1}} \right] {}_0\partial_{x_i}^{p-n} u - \sum_{j \neq i} \binom{p}{n} \frac{\mathrm{d}^n \xi^j}{\mathrm{d}\,x_i^n} \frac{\partial}{\partial x_j} \, {}_0\partial_{x_i}^{p-n} u \right\} \\
& + \sum_{n=2}^{\infty} \sum_{m=2}^{n} \sum_{k=2}^{m} \sum_{r=0}^{k-1} \binom{p}{n} \binom{n}{m} \binom{k}{r} \frac{{x_i}^{n-p} (-u)^r}{k! \Gamma(n+1-p)} \frac{\mathrm{d}^m \, u^{k-r}}{\mathrm{d}\,x_i^m} \frac{\partial^{n-m+k} \phi}{\partial x_i^{n-m} \partial u^k}.
\end{aligned} \tag{4.4}$$

After some manipulations, we get the following determining equation

$$\begin{aligned}
\frac{\partial \phi}{\partial t} = {}& - \left(\frac{\partial \phi}{\partial u} - \frac{\mathrm{d}\,\tau}{\mathrm{d}\,t} \right) \sum_{i=1}^{N} \alpha_i u^{\beta_i} \, {}_0\partial_{x_i}^{p_i} u + \sum_{i=1}^{N} \frac{\mathrm{d}\,\xi^i}{\mathrm{d}\,t} \frac{\partial u}{\partial x_i} + \\
& \phi \sum_{i=1}^{N} \alpha_i \beta_i u^{\beta_i - 1} \, {}_0\partial_{x_i}^{p_i} u + \sum_{i=1}^{N} \alpha_i u^{\beta_i} \left[{}_0\partial_{x_i}^{p_i} \phi + {}_0\partial_{x_i}^{p_i} u \left(\frac{\partial \phi}{\partial u} - p_i \frac{\mathrm{d}\,\xi^i}{\mathrm{d}\,x_i} \right) - u \, {}_0\partial_{x_i}^{p_i} \frac{\partial \phi}{\partial u} \right. \\
& + \sum_{n=1}^{\infty} \left(\binom{p_i}{n} \frac{\partial^{n+1} \phi}{\partial x_i^n \partial u} - \binom{p_i}{n+1} \frac{\mathrm{d}^{n+1} \xi^i}{\mathrm{d}\,x_i^{n+1}} \right) {}_0\partial_{x_i}^{p_i - n} u \\
& \left. - \sum_{n=1}^{\infty} \binom{p_i}{n} \left(\frac{\mathrm{d}^n \tau}{\mathrm{d}\,x_i^n} \frac{\partial}{\partial t} \, {}_0\partial_{x_i}^{p_i - n} u + \sum_{j \neq i} \frac{\mathrm{d}^n \xi^j}{\mathrm{d}\,x_i^n} \frac{\partial}{\partial x_j} \, {}_0\partial_{x_i}^{p_i - n} u \right) \right] \\
& + \sum_{i=1}^{N} \alpha_i u^{\beta_i} \sum_{n=2}^{\infty} \sum_{m=2}^{n} \sum_{k=2}^{m} \sum_{r=0}^{k-1} \binom{p}{n} \binom{n}{m} \binom{k}{r} \frac{{x_i}^{n-p} (-u)^r}{k! \Gamma(n+1-p)} \frac{\mathrm{d}^m \, u^{k-r}}{\mathrm{d}\,x_i^m} \frac{\partial^{n-m+k} \phi}{\partial x_i^{n-m} \partial u^k}.
\end{aligned} \tag{4.5}$$

The right hand side can be written as a sum over $i = 1, \dots, N$ in which the i-th addend is given by

(4.6)
$$\left[\left(\frac{\mathrm{d}\,\tau}{\mathrm{d}\,t} - \frac{\partial\phi}{\partial u}\right)\alpha_i u^{\beta_i} + \frac{\mathrm{d}\,\xi^i}{\mathrm{d}\,t}\frac{\partial u}{\partial x_i} + \phi\alpha_i\beta_i u^{\beta_i-1} + \alpha_i u^{\beta_i}\left(\frac{\partial\phi}{\partial u} - p_i\frac{\mathrm{d}\,\xi^i}{\mathrm{d}\,x_i}\right)\right]\,{}_0\partial^{p_i}_{x_i}\,u$$
$$+\,\alpha_i u^{\beta_i}\sum_{n=1}^{\infty}\left[\binom{p_i}{n}\frac{\partial^{n+1}\phi}{\partial x_i^n\partial u} - \binom{p_i}{n+1}\frac{\mathrm{d}^{n+1}\,\xi^i}{\mathrm{d}\,x_i^{n+1}}\right]\,{}_0\partial^{p_i-n}_{x_i}\,u$$
$$+\alpha_i u^{\beta_i}\left[{}_{x_i}\partial^{p_i}_0\,\phi - u\,{}_0\partial^{p_i}_{x_i}\frac{\partial\phi}{\partial u} - \sum_{n=1}^{\infty}\binom{p_i}{n}\left(\frac{\mathrm{d}^n\,\tau}{\mathrm{d}\,x_i^n}\frac{\partial}{\partial t}\,{}_0\partial^{p_i-n}_{x_i}\,u - \sum_{j\neq i}\frac{\mathrm{d}^n\,\xi^j}{\mathrm{d}\,x_i^n}\frac{\partial}{\partial x_j}\,{}_0\partial^{p_i-n}_{x_i}\,u\right)\right]$$
$$+\,\alpha_i u^{\beta_i}\sum_{n=2}^{\infty}\sum_{m=2}^{n}\sum_{k=2}^{m}\sum_{r=0}^{k-1}\binom{p}{n}\binom{n}{m}\binom{k}{r}\frac{{x_i}^{n-p}(-u)^r}{k!\Gamma(n+1-p)}\frac{\mathrm{d}^m\,u^{k-r}}{\mathrm{d}\,x_i^m}\frac{\partial^{n-m+k}\phi}{\partial x_i^{n-m}\partial u^k}.$$

From the previous equation we deduce the relations

$$\frac{n+1}{p_i-n}\frac{\partial^{n+1}\phi}{\partial x_i^n\partial u} - \frac{\mathrm{d}^{n+1}\,\xi^i}{\mathrm{d}\,x_i^{n+1}} = 0, \qquad \forall n\in\mathbb{N}, \quad i = 1,\dots,N, \tag{4.7a}$$

$$\frac{\mathrm{d}^n\,\tau}{\mathrm{d}\,x_i^n} = 0, \quad \frac{\mathrm{d}^n\,\xi^j}{\mathrm{d}\,x_i^n} = 0, \qquad i\neq j, \quad \forall i, \quad \forall n\in\mathbb{N}. \tag{4.7b}$$

Imposing the additional constraint $\xi^j\big|_{x_j=0} = 0$, $j = 1,\dots,N$, we obtain

$$\xi^i \equiv \xi^i(x_i) = c_1^i(x_0)x_i + c_2^i(x_0)x_i^2, \quad i = 1,\dots,N, \tag{4.8a}$$

$$\tau \equiv \tau(t), \tag{4.8b}$$

$$\frac{\partial^2\partial\phi}{\partial x_i\partial u} = (p_i-1)c_2^i(t). \tag{4.8c}$$

The latter relations imply $\mu^{p_i}_{x_i} = 0$, $\forall i = 1,\dots,N$. Finally we have

$$\tau(x_1) = c_1 t + c_0, \tag{4.9a}$$

$$\xi^i(x_i) = \frac{c_1 - \beta_i c_2}{p_i}x_i, \quad i = 1,\dots,N, \tag{4.9b}$$

$$\phi = c_2 u. \tag{4.9c}$$

If $\beta_i = 0\ \forall i$, denoting by $\gamma(t, x_1, \dots, x_N)$ a solution of eq. (4.1), we get $\phi = c_2 u + \gamma$.

Assuming that at least one value of β_i is different from zero, we obtain three symmetry generators,

$$\mathbf{v}_1 = \partial_t, \tag{4.10a}$$

$$\mathbf{v}_2 = t\partial_t + \sum_{i=1}^{N}\frac{x_i}{p_i}\partial_{x_i}, \tag{4.10b}$$

$$\mathbf{v}_3 = u\partial_u - \sum_{i=1}^{N}\frac{\beta_i x_i}{p_i}\partial_{x_i}. \tag{4.10c}$$

The first generator expresses the trivial invariance of equation (4.1) under translations in the variable t. The corresponding reduced equation can be obtained by looking for a stationary solution $u = v(x_1, \dots, x_N)$. The generators $\mathbf{v}_2$ and $\mathbf{v}_3$ express the invariance of the equation under scaling transformations in the dependent and independent variables.

Let us analyze in more detail the generator $\mathbf{v}_2$. By means of the equation $\mathbf{v}_2 z(x,u)=0$, we can determine N invariants of the form

$$z_i = t x_i^{-\frac{1}{p_i}}, \quad i=1,\dots,N,$$

and search for a solution of eq. (4.1) of the type

$$u(t,x_1,\dots,x_N) \equiv v\left(t^{-p_1}x_1,\dots,t^{-p_N}x_N\right). \tag{4.11}$$

The reduced equation is easily obtained in terms of the new variables $z_i = t^{-p_i}x_i$ as

$$\sum_{i=1}^{N}\left[p_i z_i \frac{\partial}{\partial z_i} + \alpha_i v^{\beta_i}\, {}_0\partial_{z_i}^{p_i}\right] v(z_1,\dots,z_N) = 0. \tag{4.12}$$

Similarly, the symmetry generator $\mathbf{v}_3$ provides the invariants

$$v = x_1^{-\frac{p_1}{\beta_1}} u, \qquad z_i = x_i x_1^{-\frac{\beta_i p_1}{\beta_1 p_i}} \quad i=2,\dots,N.$$

From them, we compute the reduced equation for $v = v(t,z_2,\dots,z_N)$:

$$\partial_t v = z_2^{\frac{\beta_2}{p_2}} \prod_{j=0}^{[p]}\left(1-p+j-\frac{1}{2}\frac{\beta_2 p_1}{\beta_1 p_2} z_2 \frac{\partial}{\partial z_2}\right) \mathfrak{K}^{[p]-p}_{\left\{\frac{\beta_i p_1}{\beta_1 p_i}\right\}}\left[z_2^{-\frac{\beta_2}{p_2}} v\right] + \sum_{i=2}^{N} \alpha_i v^{\beta_i}\, {}_0\partial_{z_i}^{p_i} v. \tag{4.13}$$

Here we used the fact that, given a function of $N+1$ variables $f(t,x_1,\dots,x_n) \equiv f\left(x_1 t^{-\alpha_1},\dots,x_n t^{-\alpha_n}\right)$, $\alpha_i > 0\ \forall i$, we can write

$${}_0\partial_t^p f = \frac{\partial^{[p]+1}}{\partial t^{[p]+1}}\left[t^{[p]-p+1}\mathfrak{K}^{[p]-p}_{\left\{\frac{1}{\alpha_i}\right\}}[f](x_1 t^{-\alpha_1},\dots,x_N t^{-\alpha_N})\right]. \tag{4.14}$$

In the previous equations, we have introduced the following integral operator:

$$\mathfrak{K}^{[p]-p}_{\left\{\frac{1}{\alpha_i}\right\}}[f](z_1,\dots,z_n) \coloneqq \int_1^{+\infty} \frac{(\omega-1)^{[p]-p}}{\Gamma([p]+1-p)} f\left(z_1\omega^{\alpha_1},\dots,z_n\omega^{\alpha_n}\right)\omega^{p-[p]-2}\,\mathrm{d}\,\omega, \tag{4.15}$$

that represents a natural generalization of the standard *Erdély–Kober fractional integral operator* [2].

Acknowledgments

G. S. acknowledges the financial support of the John Templeton Foundation. The research of P. T. has been partly supported by the research project FIS2015-63966, MINECO, Spain, and by the ICMAT Severo Ochoa project SEV-2015-0554 (MINECO).

References

[1] G.W. Bluman and S. Kumei. *Symmetries and Differential Equations*. Number v. 81 in Applied mathematical sciences. World Publishing Corporation, 1989.
[2] E. Buckwar and Y. Luchko. Invariance of a Partial Differential Equation of Fractional Order under the Lie Group of Scaling Transformations. *Journal of Mathematical Analysis and Applications*, 227(1):81–97, November 1998.
[3] R.K. Gazizov, A.A. Kasatkin, and S.Y. Lukashchuk. Symmetry properties of fractional diffusion equations. *Physica Scripta*, T136:014016, October 2009.
[4] R.K. Gazizov, A.A. Kasatkin, and S.Y. Lukashchuk. Group-Invariant solutions of Fractional Differential Equations. *Nonlinear Science and Complexity*, pages 51–59, 2011.
[5] N.H. Ibragimov. *Transformation Groups Applied to Mathematical Physics*. Mathematics and its Applications. Springer Netherlands, 2001.
[6] A.A. Kilbas, H.M. Srivastava, and J.J. Trujillo. *Theory and Applications of Fractional Differential Equations*. North-Holland mathematics studies. Elsevier, 2006.

[7] J. Krasilshchik and A. Verbovetsky. Geometry of jet spaces and integrable systems. *Journal of Geometry and Physics*, 61(9):1633–1674, 2011.
[8] R.A. Leo, G. Sicuro, and P. Tempesta. A theorem on the existence of symmetries of fractional PDEs. *Comptes Rendus Mathematique*, 352(3):219–222, 2014.
[9] K.B. Oldham and J. Spanier. *The Fractional Calculus*. Mathematics in science and engineering. Academic Press, 1974.
[10] P.J. Olver. *Applications of Lie Groups to Differential Equations*. Graduate Texts in Mathematics. Springer New York, 2000.
[11] T.J. Osler. Leibniz rule for fractional derivatives generalized and an application to infinite series. *SIAM Journal on Applied Mathematics*, 18(3):658–674, 1970.
[12] S. Samko, A.A. Kilbas, and O. Marichev. *Fractional Integrals and Derivatives*. Taylor & Francis, 1993.
[13] D.J. Saunders. *The Geometry of Jet Bundles*. Cambridge Studies in Population, Economy, and Society in Pas. Cambridge University Press, 1989.
[14] A.M. Vinogradov. Symmetries and conservation laws of partial differential equations: Basic notions and results. *Acta Applicandae Mathematicae*, 15(1-2):3–21, 1989.
[15] A.M. Vinogradov and J. Krasilshchik, editors. *Symmetries and Conservation Laws for Differential Equations of Mathematical Physics*. American Mathematical Society.

(R.A. Leo) Dipartimento di Matematica e Fisica "Ennio De Giorgi", Università del Salento, Via per Arnesano, 73100 – Lecce, Italy
E-mail address: leora@le.infn.it

(G. Sicuro) Centro Brasileiro de Pesquisas Físicas, Rua Dr. Xavier Sigaud, 150, 22290-180, Rio de Janeiro, Brazil
E-mail address: sicuro@cbpf.br

(P. Tempesta) Departamento de Física Teórica II (Métodos Matemáticos de la física), Facultad de Físicas, Universidad Complutense de Madrid, 28040 – Madrid, Spain and Instituto de Ciencias Matemáticas (CSIC-UAM-UC3M-UCM), C/ Nicolás Cabrera, No 13–15, 28049 Madrid, Spain
E-mail address: p.tempesta@fis.ucm.es,piergiulio.tempesta@icmat.es